

Tutte polynomial and G -parking functions

Hungyung Chang^a

Jun Ma^{b,*}

Yeong-Nan Yeh^{c,†}

^{a,b,c} Institute of Mathematics, Academia Sinica, Taipei, Taiwan

Abstract

Let G be a connected graph with vertex set $\{0, 1, 2, \dots, n\}$. We allow G to have multiple edges and loops. In this paper, we give a characterization of external activity by some parameters of G -parking functions. In particular, we give the definition of the bridge vertex of a G -parking function and obtain an expression of the Tutte polynomial $T_G(x, y)$ of G in terms of G -parking functions. We find the Tutte polynomial enumerates the G -parking function by the number of the bridge vertices.

Keywords: parking functions; spanning tree; Tutte polynomial

1 Introduction

J. Riordan [12] define the parking function as follows: m parking spaces are arranged in a line, numbered 1 to n left to right; n cars, arriving successively, have initial parking preferences, a_i for i , chosen independently and at random; $(a_1, \dots, a_n)$ is called preference function; if space a_i is occupied, car i moves to the first unoccupied space to the right; if all the cars can be parked, then the preference function is called parking function.

Konheim and Weiss [8] introduced the conception of the parking functions of length n in the study of the linear probes of random hashing function. J. Riordan [12] studied the parking functions and derived that the number of parking functions of length n is $(n+1)^{n-1}$, which coincides with the number of labeled trees on $n+1$ vertices by Cayley's formula. Several bijections between the two

*Email address of the corresponding author: majun@math.sinica.edu.tw

†Partially supported by NSC 96-2115-M-001-005

sets are known (e.g., see [5, 12, 13]). Parking functions have been found in connection to many other combinatorial structures such as acyclic mappings, polytopes, non-crossing partitions, non-nesting partitions, hyperplane arrangements, etc. Refer to [6, 5, 7, 11, 14, 15] for more information.

Parking function $(a_1, \dots, a_n)$ can be redefined that its increasing rearrangement $(b_1, \dots, b_n)$ satisfies $b_i \leq i$. Pitman and Stanley generalized the notion of parking functions in [11]. Let $\mathbf{x} = (x_1, \dots, x_n)$ be a sequence of positive integers. The sequence $\alpha = (a_1, \dots, a_n)$ is called an $\mathbf{x}$ -parking function if the non-decreasing rearrangement $(b_1, \dots, b_n)$ of α satisfies $b_i \leq x_1 + \dots + x_i$ for any $1 \leq i \leq n$. Thus, the ordinary parking function is the case $\mathbf{x} = (1, \dots, 1)$. By the determinant formula of Gončarov polynomials, Kung and Yan [9] obtained the number of $\mathbf{x}$ -parking functions for an arbitrary $\mathbf{x}$. See also [16, 17, 18] for the explicit formulas and properties for some specified cases of $\mathbf{x}$.

Recently, Postnikov and Shapiro [10] gave a new generalization, building on work of Cori, Rossin and Salvy [1], the G -parking functions of a graph. For the complete graph $G = K_{n+1}$, the defined functions in [10] are exactly the classical parking functions. Chebikin and Pylyavskyy [2] established a family of bijections from the set of G -parking functions to the spanning trees of G .

Dimitrije Kostic and Catherine H. Yan [3] proposed the notion of a G -multiparking function, a natural extension of the notion of a G -parking function and extended the result of [18] to arbitrary graphs. They constructed a family of bijections from the set of G -multiparking functions to the spanning forests of G . Particularly, They characterize the external activity by the bijection induced by the breadth-first search and gave a representation of Tutte polynomial by the reversed sum of G -multiparking functions. Given a classical parking function $\alpha = (a_1, \dots, a_n)$, let $cr(\alpha)$ be the number of critical left-to-right maxima in α . They also gave an expression of the Tutte polynomial $T_{K_{n+1}}(x, y)$ of the complete graph K_{n+1} as follows:

$$T_{K_{n+1}}(x, y) = \sum_{\alpha \in \mathcal{P}_n} x^{cr(\alpha)} y^{\binom{n}{2} - \sum_{i=1}^n a_i},$$

where $\mathcal{P}_n$ is the set of classical parking functions of length n . Recently, Sen-peng Eu, Tung-Shan Fu and Chun-Ju Lai [4] considered a class of multigraphs in connection with $\mathbf{x}$ -parking functions, where $\mathbf{x} = (a, b, \dots, b)$. They gave the Tutte polynomial of the multigraphs in terms of $\mathbf{x}$ -parking

functions.

Let G be a connected graph with vertex set $\{0, 1, 2, \dots, n\}$. We allow G to have multiple edges and loops. The motivation of this paper is to extend the results in [18] on K_{n+1} to arbitrary connected graphs and give a characterization of external activity by some parameters of G -parking functions. To obtain the characterization for the complete graph K_{n+1} , Dimitrije Kostic and Catherine H. Yan [3] use the bijections induced by the breadth-first search. In this paper, we use the bijections induced by the vertex ranking. We give the definition of the bridge vertex of a G -parking functions. We obtain an expression of the Tutte polynomial $T_G(x, y)$ of G in terms of G -parking functions. So, we find the Tutte polynomial enumerates the G -parking function by the number of the bridge vertices.

This paper is organized as follows. In Section 2, we give the definition of the bridge vertex of a G -parking functions. In Section 3, we will express the Tutte polynomial $T_G(x, y)$ of G in terms of G -parking functions.

2 The bridge vertex of G -parking function

In this section, we always let G be a connected graph with vertex set $\{0, 1, 2, \dots, n\}$ and edge set $E(G)$. We allow G to have multiple edges and loops. Let $[n] := \{1, 2, \dots, n\}$. For any $I \subseteq V(G) \setminus \{0\}$ and $v \in I$, define $\text{outdeg}_{I,G}(v)$ to be the cardinality of the set $\{\{w, v\} \in E(G) \mid w \notin I\}$. We give the definition of G -parking function as follows.

Definition 2.1 *Let G be a connected graph with vertex set $V(G) = \{0, 1, 2, \dots, n\}$ and edge set $E(G)$. A G -parking function is a function $f : V(G) \rightarrow \mathbb{N} \cup \{-1\}$, such that for every $I \subseteq V(G) \setminus \{0\}$ there exists a vertex $v \in I$ such that $0 \leq f(v) < \text{outdeg}_{I,G}(v)$ and $f(0) = -1$.*

For any $i, j \in [n]$, let $\mu_G(i, j)$ be the number of edges connecting the vertices i to j in G . For establishing the bijections, all edges of G are colored. The colors of edges connecting the vertices i to j are $0, 1, \dots, \mu_G(i, j) - 1$ respectively for any $i, j \in V(G)$. We use $\{i, j\}_k$ to denote the edge $e \in E(G)$ connecting two vertices i and j with color k . A subgraph T of G is called a subtree of G

rooted at m if the subgraph contains the vertex m and there is a unique path from i to m in T for every vertex i of T . If a subtree contains all vertices of G , then we say the subtree is a spanning tree of G . Let $\mathcal{P}_G$ and $\mathcal{T}_G$ be the sets of the G -parking functions and the color spanning trees of G respectively. For any $T \in \mathcal{T}_G$ and $e \in T$, let $c_T(e)$ denote the color of edge e in T . Kostic and Yan [3] give an algorithm Φ which is a bijection from the sets $\mathcal{P}_G$ to $\mathcal{T}_G$. We give a description of the algorithm as follows.

Algorithm A. (Kostic, Yan [3])

Step 1: Let $val_0 = f$, $P_0 = \emptyset$, $T_0 = Q_0 = \{0\}$.

Step 2: At time $i \geq 1$, let $v = \min\{\tau(w) \mid w \in Q_{i-1}\}$, where τ is a vertex ranking in S_n .

Step 3: Let $N = \{w \notin P_{i-1} \mid 0 \leq val_{i-1}(w) \leq \mu(w, v) - 1 \text{ and } \{w, v\}_{val_{i-1}(w)} \in E(G)\}$ and $\hat{N} = \{w \notin P_{i-1} \mid val_{i-1}(w) \geq \mu(w, v) \text{ and } \{w, v\}_{val_{i-1}(w)} \in E(G)\}$. Set $val_i(w) = val_{i-1}(w) - \mu_G(w, v)$ for all $w \in \hat{N}$. For any other vertex u , set $val_i(u) = val_{i-1}(u)$. Update P_i , Q_i and F_i by letting $P_i = P_{i-1} \cup \{v\}$, $Q_i = Q_{i-1} \cup N \setminus \{v\}$. Let T_i be a graph on $P_i \cup Q_i$ whose edges are obtained from those of T_{i-1} by joining edges $\{w, v\}_{val_{i-1}(w)}$ for each $w \in N$.

Define $\Phi = \Phi_{G, \tau} : \mathcal{P}_G \rightarrow \mathcal{T}_G$, by letting $\Phi(f) = T_n$. Let $T \in \mathcal{F}_G$. Note that the vertex 0 is the root of T . For any non-root vertex $v \in [n]$, there is a unique path from v to 0 in T . Define the height of v to be the number of edges in the path. If the height of a vertex w is less than the height of v and $\{v, w\}_k$ is an edge of T , then w is the predecessor of v , v is a child of w , and write $w = pre_T(v)$ and $v \in child_T(w)$. Suppose T' is a subtree of G . A leaf of T' is a vertex of T' with degree 1 in T' . Denote the set of leaves of T' by $Leaf(T')$. The following algorithm will give the inverse map of Φ .

Algorithm B (Kostic, Yan [3]).

Step 1. Let τ be a vertex ranking in S_n . Assume $v_0, v_1, v_2, \dots, v_i$ are determined, where $v_0 = 0$. Let $V_i = \{v_0, v_1, v_2, \dots, v_i\}$ and $W_i = \{v \notin V_i \mid \{v, w\}_k \in T \text{ for some } w \in V_i\}$. Let T' be

the subtree obtained by restricting T to $V_i \cup W_i$. Let v_{i+1} be the vertex w such that $\tau(w) \leq \tau(u)$ for all $u \in \text{Leaf}(T')$.

Step 2. Let $\pi = (v_1 \dots v_n)$ be the order of the vertices of G determined by Step 1. Set $f(0) = -1$. For any other vertex v , let $f(v)$ be equal to the sum of the color of the edge connecting the vertices v and $\text{pre}_T(v)$ and the cardinality of the set $N(v)$, where $N(v) = N_{G,T,\tau}(v) = \{w \mid \{v, w\}_k \in E(G) \text{ and } \pi^{-1}(w) < \pi^{-1}(\text{pre}_T(v))\}$.

Define $\Theta = \Theta_{G,\tau} : \mathcal{T}_G \rightarrow \mathcal{P}_G$ by letting $\Theta_{G,\tau}(T) = f_T$. Then Θ is the inverse of Φ . Note that the order $\pi = v_1 v_2 \dots v_n$ in the algorithm B is exactly the order in which vertices of G will be placed into the set P_i when running algorithm A on f . Define $\text{Ord} = \text{Ord}_{G,\tau} : \mathcal{P}_G \rightarrow S_n$ by letting $\text{Ord}_{G,\tau}(f) = (v_1 v_2 \dots v_n)$, where the order $0 = v_0, v_1, v_2, \dots v_n$ is obtained by Algorithm B, i.e., $\text{Ord}_{G,\tau}(f)_i = u$ and $\text{Ord}_{G,\tau}^{-1}(f)_u = i$ if $v_i = u$ for all $i \in [n]$. Furthermore, let $\text{Rea}(f)$ be a function such that $\text{Rea}(f)(i) = f(\text{Ord}(f)_i)$ for all $i \in [n]$. We say $\text{Ord}(f)$ and $\text{Rea}(f)$ are an order and a rearrangement of the G -parking function f respectively. Hence, for any $f \in \mathcal{P}_G$, we can obtain a pair $(\text{Rea}(f), \text{Ord}(f))$.

Definition 2.2 Let $f \in \mathcal{P}_G$ and $v \in V(G)$. Suppose $\text{Ord}(f)_i = v$. let $I_v = I_{G,\tau,f,v} = \{\text{Ord}(f)_j \mid j \geq i\}$. The vertex v is said to be f -critical if $f(v) = \text{outdeg}_{I_v}(v) - 1$.

Define $C_f = C_{G,\tau,f}$ to be the set of all the f -critical vertices. Clearly, $C_f \neq \emptyset$ for any $f \in \mathcal{P}_G$ since $0 \in C_f$.

Example 2.3 Let us consider the following graph G . Let τ be the identity permutation.




Fig 1. A graph G

We list all the G -parking functions f as well as the corresponding $\text{Ord}(f)$, $\text{Rea}(f)$ and C_f as follows.

G -parking functions f	$\text{Ord}(f)$	$\text{Rea}f$	C_f
$f_1 = (-1, 0, 0, 0)$	$(0, 1, 2, 3)$	$(-1, 0, 0, 0)$	$\{0, 1, 2\}$
$f_2 = (-1, 0, 0, 1)$	$(0, 1, 2, 3)$	$(-1, 0, 0, 1)$	$\{0, 1, 2\}$
$f_3 = (-1, 0, 0, 2)$	$(0, 1, 2, 3)$	$(-1, 0, 0, 2)$	$\{0, 1, 2, 3\}$
$f_4 = (-1, 0, 1, 0)$	$(0, 1, 3, 2)$	$(-1, 0, 0, 1)$	$\{0, 1, 2\}$
$f_5 = (-1, 0, 1, 1)$	$(0, 1, 3, 2)$	$(-1, 0, 1, 1)$	$\{0, 1, 2, 3\}$
$f_6 = (-1, 1, 0, 0)$	$(0, 3, 1, 2)$	$(-1, 0, 1, 0)$	$\{0, 1, 3\}$
$f_7 = (-1, 1, 1, 0)$	$(0, 3, 1, 2)$	$(-1, 0, 1, 1)$	$\{0, 1, 2, 3\}$
$f_8 = (-1, 2, 0, 0)$	$(0, 3, 2, 1)$	$(-1, 0, 0, 2)$	$\{0, 1, 2, 3\}$

Table 1. All the G -parking functions

Definition 2.4 Let $f \in \mathcal{P}_G$ and $v \in V(G) \setminus \{0\}$. Suppose $\text{Ord}(f)_i = v$. A G -parking function g is weak v -identical to f if it satisfies the following conditions:

- (1) $\text{Rea}(g)(j) = \text{Rea}(f)(j)$ and $\text{Ord}(g)_j = \text{Ord}(f)_j$ for all $j \in [i - 1]$,
- (2) $g(v) \geq f(v)$, and
- (3) $g(w) \geq \text{outdeg}_{I_v}(w)$ for all $w \in I_v$ and $\tau(w) < \tau(v)$.

Furthermore, g is strong v -identical to f if (1) g is weak v -identical to f ; (2) $\text{Ord}(g)_i = v$.

Given $f \in \mathcal{P}_G$ and $v \in V(G) \setminus \{0\}$, define

$$W_{v,f} = W_{G,v,\tau,f} = \{g \in \mathcal{P}_G \mid g \text{ is weak } v\text{-identical to } f\}$$

and

$$S_{v,f} = S_{G,v,\tau,f} = \{g \in \mathcal{P}_G \mid g \text{ is strong } v\text{-identical to } f\}.$$

It is easy to see that $S_{v,f} \subseteq W_{v,f}$ and $g(v) = f(v)$ for all $g \in S_{v,f}$ if $v \in C_f$.

Lemma 2.5 Let G be a connected graph and f a G -parking function. Let e be an edge of G connecting the vertices w to v . Suppose that e is a bridge of G and the vertices w and 0 are in the same component after deleting the edge e . Then $v \in B(f)$ for any $f \in \mathcal{P}_G$.

Proof. Since e is a bridge of G and the vertices w and 0 are in the same component after deleting the edge e , we have $f(v) = 0$, $\text{Ord}(f)^{-1}(w) < \text{Ord}(f)^{-1}(v)$ and $v \in C_f$ for all $f \in \mathcal{P}_G$. Given a $f \in \mathcal{P}_G$, suppose $\text{Ord}(f)_i = v$. Assume that $W_{v,f} \neq S_{v,f}$, i.e., there is a $g \in \mathcal{P}_G$ such that $g \in W_{v,f}$ and $g \notin S_{v,f}$. $g \notin S_{v,f}$ implies $\text{Ord}(g)_i \neq v$. Let $u = \text{Ord}(g)_i$. Then $\tau(u) > \tau(v)$ and $\text{Ord}(g)^{-1}(u) < \text{Ord}(g)^{-1}(v)$ since $g \in W_{v,f}$. Note that $\text{Ord}(g)^{-1}(w) = \text{Ord}(f)^{-1}(w) < \text{Ord}(f)^{-1}(v) = \text{Ord}(g)^{-1}(u)$. So, we must have $\text{Ord}(g)^{-1}(v) < \text{Ord}(g)^{-1}(u)$ by Algorithm A, a contradiction. Hence, $W_{v,f} = S_{v,f}$ and $v \in B(f)$. $\blacksquare$

Definition 2.6 Let $f \in \mathcal{P}_G$. A vertex $v \in V(G) \setminus \{0\}$ is said to be f -bridge if $v \in C_f$ and $|W_{v,f}| = |S_{v,f}|$.

Define $B(f) = B_{G,\tau}(f)$ as the set of the f -bridge vertices of f , $b(f) = b_{G,\tau}(f) = |B_{G,\tau}(f)|$ and $w(f) = w_G(f) = |E(G)| - |V(G)| - \sum_{i=0}^n f(i)$.

Example 2.7 We consider the graph G in Fig 1. By Table 1, we have $f_3 = (-1, 0, 0, 2)$ is a G -parking function and $C_{f_3} = \{0, 1, 2, 3\}$. It is easy to check the results in Table 2.

$W_{1,f_3} = \{f_i \mid i \in [8]\}$	$S_{1,f_3} = \{f_i \mid i \in [5]\}$
$W_{2,f_3} = \{f_i \mid i \in [5]\}$	$S_{2,f_3} = \{f_i \mid i \in [3]\}$
$W_{3,f_3} = \{f_i \mid i \in [3]\}$	$S_{3,f_3} = \{f_i \mid i \in [3]\}$

Table 2. W_{v,f_3} and S_{v,f_3} , where $v \in C_{f_3}$

Hence, $B(f_3) = \{3\}$. We list all the G -parking functions as well as the corresponding parameters

$b(f)$ and $w(f)$ in the following table.

G -parking function f	$B(f)$	$(b(f), w(f))$
$f_1 = (-1, 0, 0, 0)$	$\emptyset$	$(0, 2)$
$f_2 = (-1, 0, 0, 1)$	$\emptyset$	$(0, 1)$
$f_3 = (-1, 0, 0, 2)$	$\{3\}$	$(1, 0)$
$f_4 = (-1, 0, 1, 0)$	$\{2\}$	$(1, 1)$
$f_5 = (-1, 0, 1, 1)$	$\{2, 3\}$	$(2, 0)$
$f_6 = (-1, 1, 0, 0)$	$\{3\}$	$(1, 1)$
$f_7 = (-1, 1, 1, 0)$	$\{2, 3\}$	$(2, 0)$
$f_8 = (-1, 2, 0, 0)$	$\{1, 2, 3\}$	$(3, 0)$

Table 3. G -parking functions f as well as the corresponding parameters $b(f)$ and $w(f)$

We note that the Tutte polynomial $T_G(x, y)$ of G in Fig.1 satisfies

$$T_G(x, y) = x^3 + 2x^2 + x + 2xy + y + y^2 = \sum_{f \in \mathcal{P}_G} x^{b(f)} y^{w(f)}.$$

3 A new expression of the Tutte polynomial

In this section, we will prove the main theorem of this paper. Suppose that e is an edge connecting the vertices i to j in G , where $i < j$. Define a graph $G \setminus e$ as follows. The graph $G \setminus e$ is obtained from G contracting the the vertices i and j ; that is, to get $G \setminus e$ we identify two vertices i and j as a new vertex i . Define $G - e$ as a graph obtained by deleting the edge e from G .

Let $NB_G(0)$ be the set of the vertices which are adjacent to the vertex 0 in G . We consider the case in which e is an edge of G connecting the root 0 to the vertex u and u satisfies $\tau(u) \leq \tau(w)$ for all $w \in NB_G(0)$. Let $\mathcal{P}_G^0 = \{f \in \mathcal{P}_G \mid f(u) = 0\}$ and $\mathcal{P}_G^1 = \{f \in \mathcal{P}_G \mid f(u) \geq 1\}$. Clearly, $\mathcal{P}_G^0 \cap \mathcal{P}_G^1 = \emptyset$ and $\mathcal{P}_G = \mathcal{P}_G^0 \cup \mathcal{P}_G^1$. For any $f \in \mathcal{P}_G^0$, let $g = \phi(f)$ such that $g(w) = f(w)$ for any $w \neq u$. For any $f \in \mathcal{P}_G^1$, let $g = \varphi(f)$ such that $g(w) = f(w)$ for any $w \neq u$ and $g(u) = f(u) - 1$.

Lemma 3.1 (1) The mapping ϕ is a bijection from $\mathcal{P}_G^0$ to $\mathcal{P}_{G \setminus e}$ with $w_{G \setminus e}(\phi(f)) = w_G(f)$.

(2) For any $f \in \mathcal{P}_G^0$, we have $B_{G, \tau}(f) \setminus \{u\} = B_{G/e, \tau}(\phi(f))$.

Proof. (1) For any $I \subset V(G \setminus e)$ with $0 \notin I$ and $w \in I$, we have $\text{outdeg}_{I, G \setminus e}(w) = \text{outdeg}_{I, G}(w)$. This implies $g = \phi(f)$ is a $(G \setminus e)$ -parking function. Conversely, for any $g \in \mathcal{P}_{G \setminus e}$, let $f = \phi^{-1}(g)$ such that $f(w) = g(w)$ for any $w \in V(G \setminus e)$ and $f(u) = 0$. For any $I \subset V(G)$ with $0 \notin I$, if $u \in I$, then $f(u) < \text{outdeg}_{I, G}(u)$ since $f(u) = 0$ and $\text{outdeg}_{I, G}(u) \geq 1$; otherwise, we have $\text{outdeg}_{I, G \setminus e}(w) = \text{outdeg}_{I, G}(w)$ for all $w \in I$, this implies $f(w) < \text{outdeg}_{I, G}(w)$ for some $w \in I$ since g is a $(G \setminus e)$ -parking function. Clearly, $w_{G \setminus e}(g) = w_G(f)$.

(2) Since $\tau(u) < \tau(w)$ for all $w \in NB_G(0) \setminus \{u\}$ and $f(u) = 0$, we have $\text{Ord}_{G, \tau}(f)_1 = u$ for all $f \in \mathcal{P}_G^0$. Let $g = \phi(f)$. Then $\text{Ord}_{G \setminus e, \tau}(g)_i = \text{Ord}_{G, \tau}(f)_{i+1}$ for all $i \in [n-1]$. Let $v \in B_{G, \tau}(f) \setminus \{u\}$. Clearly, $\text{outdeg}_{I_v, G \setminus e}(v) = \text{outdeg}_{I_v, G}(v)$ and $f(v) = g(v)$. This implies $g(v) = \text{outdeg}_{I_v, G \setminus e}(v) - 1$. Hence, v is g -critical in $G \setminus e$ since v is f -critical in G .

Now, we assume that $W_{G \setminus e, v, \tau, g} \neq S_{G \setminus e, v, \tau, g}$, i.e., there is a $g_1 \in \mathcal{P}_{G \setminus e}$ such that $g_1 \in W_{G \setminus e, v, \tau, g}$ and $g_1 \notin S_{G \setminus e, v, \tau, g}$. Suppose $\text{Ord}_{G, \tau}(f)_i = v$. So, $\text{Ord}_{G \setminus e, \tau}(g)_i \neq v$. Furthermore, we have $\tau(\text{Ord}_{G \setminus e, \tau}(g)_i) > \tau(v)$. Let $f_1 = \phi^{-1}(g_1)$. It is easy to see that $f_1 \in W_{G, v, \tau, f}$ and $f_1 \notin S_{G, v, \tau, f}$, a contradiction.

Conversely, let $v \in B_{G \setminus e, \tau}(g)$. Clearly, $v \neq u$. Assume $W_{G, v, \tau, f} \neq S_{G, v, \tau, f}$, where $f = \phi^{-1}(g)$, i.e., there is a $f_1 \in \mathcal{P}_G$ such that $f_1 \in W_{G, v, \tau, f}$ and $f_1 \notin S_{G, v, \tau, f}$. Let $g_1 = \phi(f_1)$. Similarly, we can obtain $W_{G \setminus e, v, \tau, g} \neq S_{G \setminus e, v, \tau, g}$, a contradiction. $\blacksquare$

Lemma 3.2 (1) The mapping φ is a bijection from $\mathcal{P}_G^1$ to $\mathcal{P}_{G-e}$ with $w_{G-e}(\varphi(f)) = w_G(f)$.

(2) For any $f \in \mathcal{P}_G^1$, we have $B_{G, \tau}(f) = B_{G \setminus e, \tau}(\varphi(f))$.

Proof. Since $f(u) \geq 1$, we have the edge $\{0, u\}$ isn't a bridge. So, $G - e$ is still a connected graph. (1) For any $I \subset V(G - e)$ with $0 \notin I$ and $w \in I$, we have $\text{outdeg}_{I, G-e}(w) = \text{outdeg}_{I, G}(w)$ if $w \neq u$; $\text{outdeg}_{I, G \setminus e}(w) = \text{outdeg}_{I, G}(w) - 1$ if $w = u$. Note that $g(u) = f(u) - 1$. Hence, $g = \varphi(f)$ is a $(G - e)$ -parking function. Conversely, for any $g \in \mathcal{P}_{G-e}$, let $f = \varphi^{-1}(g)$ such that $f(w) = g(w)$ for any $w \in V(G - e)$ and $f(u) = g(u) + 1$. For any $I \subset V(G)$ with $0 \notin I$, we have $\text{outdeg}_{I, G}(w) = \text{outdeg}_{I, G-e}(w)$ if $w \neq u$; $\text{outdeg}_{I, G}(w) = \text{outdeg}_{I, G-e}(w) + 1$ if $w = u$. Note that $f(u) = g(u) + 1$. This implies $f(w) < \text{outdeg}_{I, G}(w)$ for some $w \in I$ since g is a $(G - e)$ -parking function. Clearly, $w_{G-e}(g) = w_G(f)$.

(2) Note that $\text{Ord}_{G,\tau}(f) = \text{Ord}_{G-e,\tau}(\varphi(f))$ for all $f \in \mathcal{P}_G$. For any $f \in \mathcal{P}_G$, it is easy to see that the vertex v is $\varphi(f)$ -critical in $G - e$ if and only if it is f -critical in G .

Now, given $f \in \mathcal{P}_G$, let $g = \varphi(f)$. For any $v \in B_{G,\tau}(f)$, we assume that $W_{G-e,v,\tau,g} \neq S_{G-e,v,\tau,g}$, i.e., there is a $g_1 \in W_{G-e,v,\tau,g}$ and $g_1 \notin S_{G-e,v,\tau,g}$. Suppose $\text{Ord}_{G-e,\tau}(g)_i = v$. Then $\text{Ord}_{G-e,\tau}(g_1)_i \neq v$. Furthermore, we have $\tau(\text{Ord}_{G-e,\tau}(g_1)_i) > \tau(v)$. Let $f_1 = \varphi^{-1}(g_1)$. It is easy to see that $f_1 \in W_{G,v,\tau,g}$ and $f_1 \notin S_{G,v,\tau,g}$, a contradiction.

Conversely, let $v \in B_{G-e,\tau}(g)$. Assume $W_{G,v,\tau,f} \neq S_{G,v,\tau,f}$, i.e., there is a $f_1 \in \mathcal{P}_G$ such that $f_1 \in W_{G,v,\tau,f}$ and $f_1 \notin S_{G,v,\tau,f}$. Let $g_1 = \phi(f_1)$. Similarly, we can obtain $W_{G-e,v,\tau,g} \neq S_{G-e,v,\tau,g}$, a contradiction. ■

We are in a position to prove the main theorem.

Theorem 3.3 *Suppose that G is a connected graph with vertex $\{0, 1, \dots, n\}$ and τ is a vertex ranking in S_n . Let $T_G(x, y)$ be the Tutte polynomial of G . Then $T_G(x, y) = \sum_{f \in \mathcal{P}} x^{b(f)} y^{w(f)}$.*

Proof. Let $P_G(x, y) = \sum_{f \in \mathcal{P}} x^{b(f)} y^{w(f)}$. Let e be an edge of G connecting the vertices u to 0 , where u satisfies $\tau(u) \leq \tau(w)$ for all $w \in NB_G(0)$. We consider the following three cases.

Case 1. e is a loop of G .

For any $f \in \mathcal{P}_G$, it is easy to see that f is $(G - e)$ -parking function as well. Note that

$$\begin{aligned} w_{G-e}(f) &= |E(G - e)| - |V(G - e)| - \sum_{i=0}^n f(i) \\ &= |E(G)| - 1 - |V(G)| - \sum_{i=0}^n f(i) \\ &= w_G(f) - 1. \end{aligned}$$

Hence,

$$\begin{aligned} P_G(x, y) &= \sum_{f \in \mathcal{P}_G} x^{b(f)} y^{w(f)} \\ &= \sum_{g \in \mathcal{P}_{G-e}} x^{b_\tau(g)} y^{w(g)+1} \\ &= y P_{G-e}(x, y) \end{aligned}$$

Case 2. e is a bridge of G .

For any $f \in \mathcal{P}_G$, we have $f(u) = 0$ since e is a bridge. So, $\mathcal{P}_G = \mathcal{P}_G^0$. Let ϕ be defined as that in Lemma 3.1. From Lemma 2.5, we have $u \in B_{G,\tau}(f)$ for all $f \in \mathcal{P}_G$. Lemma 3.1 (2) tells us that $b_G(f) = b_{G \setminus e}(\phi(f)) + 1$.

$$\begin{aligned} P_G(x, y) &= \sum_{f \in \mathcal{P}_G} x^{b(f)} y^{w(f)} \\ &= \sum_{g \in \mathcal{P}_{G \setminus e}} x^{b(g)+1} y^{w(g)} \\ &= x P_{G \setminus e}(x, y) \end{aligned}$$

Case 3. e is neither loop nor bridge of G .

First, we claim $u \notin B_G(f)$ for any $f \in \mathcal{P}_G^0$. Since $\tau(u) \leq \tau(w)$ for all $w \in NB_G(0)$ and $f(u) = 0$, we have $\text{Ord}_{G,\tau}(f)_1 = u$ for all $f \in \mathcal{P}_G^0$. If there are at least two edges connecting the vertices 0 and u in G , i.e., $\mu_G(0, u) \geq 2$, then u isn't f -critical for any $f \in \mathcal{P}_G^0$. So, we suppose $\mu_G(0, u) = 1$. This implies that u is f -critical for any $f \in \mathcal{P}_G^0$. Since e is neither loop nor bridge of G , there exists a $f' \in \mathcal{P}_G$ such that $f'(u) \geq 1$. Suppose $\text{Ord}_{G,\tau}(f')_1 = w$. Then $w \neq u$, $f'(w) = 0$ and $\tau(w) > \tau(u)$. Hence, $f' \in W_{G,u,\tau,f}$ and $f' \notin S_{G,u,\tau,f}$. This tells us $u \notin B_G(f)$.

By Lemmas 3.1 and 3.2, we have

$$\begin{aligned} P_G(x, y) &= \sum_{f \in \mathcal{P}_G} x^{b(f)} y^{w(f)} \\ &= \sum_{f \in \mathcal{P}_G^0} x^{b(f)} y^{w(f)} + \sum_{f \in \mathcal{P}_G^1} x^{b(f)} y^{w(f)} \\ &= \sum_{g \in \mathcal{P}_{G \setminus e}} x^{b(g)} y^{w(g)} + \sum_{g \in \mathcal{P}_{G-e}} x^{b(g)} y^{w(g)} \\ &= P_{G \setminus e}(x, y) + P_{G-e}(x, y) \end{aligned}$$

Finally, we consider the initial conditions. Let G be a graph with vertex set $\{0\}$ and $E(G) = \emptyset$. There is a unique G -parking function $f(0) = -1$. Clearly, $w(f) = 0$ and $B_G(f) = \emptyset$. So, $T_G(x, y) = 1$. Next, let G be a graph with vertex set $\{0, 1\}$ and $E(G) = \{\{0, 1\}\}$. There is a unique G -parking function $f(0) = -1$ and $f(1) = 0$. It is easy to see $B_G(f) = \{1\}$ and $w(f) = 0$. Hence, $P_G(x, y) = x$. This complete the proof. ■

Let us define a multiset $BW_G = BW_{G,\tau}$ as $BW_{G,\tau} = \{(b_{G,\tau}(f), w_G(f)) \mid f \in \mathcal{P}_G\}$. By Theorem 3.3, we immediately obtain the following corollary.

Corollary 3.4 *Let G be a connected graph. Suppose τ_1 and τ_2 are two vertex ranking. Then $BW_{G,\tau_1} = BW_{G,\tau_2}$.*

Next, we consider the case in which G is the complete graph K_{n+1} and τ is the identity permutation. Recall that the K_{n+1} -parking functions are exactly the classical parking functions, i.e., $f = (f(0), f(1), \dots, f(n)) \in \mathcal{P}_{K_{n+1}}$ if and only if $(f(1), \dots, f(n))$ is a classical parking function.

Definition 3.5 *Given a classical parking function $\alpha = (a_1, a_2, \dots, a_n)$, we say that a term $a_i = j$ is α -critical maxima if a_i satisfies that there are exactly $n - 1 - j$ terms larger than j and $k < i$ for all $a_k > j$.*

Lemma 3.6 *Let $\alpha = (a_1, a_2, \dots, a_n)$ be a classical parking function and $a_i = j$ an α -critical maxima. Then there are exactly j terms less than j .*

Proof. Let $\bar{\alpha} = (-1, a_1, a_2, \dots, a_n)$. Then $\bar{\alpha} \in \mathcal{P}_{K_{n+1}}$. Let $T = \Phi(\bar{\alpha})$ be the spanning tree obtained by Algorithm A and $v = \text{pre}_T(i)$. Then $\text{Ord}^{-1}(\bar{\alpha})_v = j$. Thus, $\text{Ord}^{-1}(\bar{\alpha})_i = j + 1$ since $a_i = j$ is an α -critical maxima. From Algorithm B, $a_{\text{Ord}(\bar{\alpha})_k} < j$ for all $1 \leq k \leq j$. ■

Lemma 3.7 *Let $\alpha = (a_1, a_2, \dots, a_n)$ be a classical parking function and $\bar{\alpha} = (-1, a_1, a_2, \dots, a_n)$. Then $a_i = j$ is α -critical maxima if and only if the vertex i is $\bar{\alpha}$ -bridge.*

Proof. First, we suppose $a_i = j$ is α -critical maxima. By Algorithm A, it is easy to see the vertex i is $\bar{\alpha}$ -critical since there are exactly j terms less than j and exactly $n - 1 - j$ terms larger than j . $k < i$ for all $a_k > a_i$ imply $W_{i,\bar{\alpha}} = S_{i,\bar{\alpha}}$. Hence, the vertex i is $\bar{\alpha}$ -bridge.

Conversely, we suppose the vertex i is $\bar{\alpha}$ -bridge, $a_i = j$ and $\text{Ord}^{-1}(\bar{\alpha})_i = j + 1$. Let $T = \Phi(\bar{\alpha})$ be the spanning tree obtained by Algorithm A and $v = \text{pre}_T(i)$. There are exactly j terms less than j and $\text{Ord}^{-1}(\bar{\alpha})_v + 1 = \text{Ord}^{-1}(\bar{\alpha})_i$ since the vertex i is $\bar{\alpha}$ -critical. Let w be the first vertex at the right of i such that $w > i$ in the order $\text{Ord}(\bar{\alpha})$. Let T' be a new spanning tree from T by deleting the edge

$\{v, i\}$ from T and adding the edge $\{i, w\}$ into T If $a_w \leq j$; otherwise adding the edge $\{v, w\}$ into T . Let $\bar{\beta} = \Theta(T')$ be the K_{n+1} -parking function by Algorithm B. Then $\bar{\beta} \in W_{i, \bar{\alpha}}$ and $\bar{\beta} \notin S_{i, \bar{\alpha}}$, a contradiction. So, we prove that for any vertex $w \in [n]$ if w is at the right of i in the order $Org(\bar{\alpha})$, then $w < i$ and $a_w > j$. Hence, a_i is α -critical maxima. ■

Let $cm(\alpha)$ be the number of critical maxima in a classical parking function α .

Corollary 3.8

$$T_{K_{n+1}}(x, y) = \sum_{\alpha \in \mathcal{P}_n} x^{cm(\alpha)} y^{\binom{n}{2} - \sum_{i=1}^n a_i}$$

where $\mathcal{P}_n$ is the set of classical parking function of length n .

Example 3.9 We list all classical the parkin functions α of length 3 as well as the corresponding sets of critical maxima and $cm(\alpha)$ in the following table.

<i>parking function</i>	<i>critical maxima</i>	$cm(\alpha)$	<i>parking function</i>	<i>critical maxima</i>	$cm(\alpha)$
(0, 0, 0)	$\emptyset$	0	(0, 0, 1)	$\emptyset$	0
(0, 0, 2)	$\{a_3\}$	1	(0, 1, 0)	$\emptyset$	0
(0, 1, 1)	$\emptyset$	0	(0, 1, 2)	$\{a_3\}$	1
(0, 2, 0)	$\{a_2\}$	1	(0, 2, 1)	$\{a_2, a_3\}$	2
(1, 0, 0)	$\emptyset$	0	(1, 0, 1)	$\emptyset$	0
(1, 0, 2)	$\{a_3\}$	1	(1, 1, 0)	$\{a_3\}$	1
(1, 2, 0)	$\{a_2, a_3\}$	2	(2, 0, 0)	$\{a_1\}$	1
(2, 0, 1)	$\{a_1, a_3\}$	2	(2, 1, 0)	$\{a_1, a_2, a_3\}$	3

Table 4. Classical the parking functions α of length n as well as their sets of critical maxima

Hence, $T_{K_4}(x, y) = y^3 + 3y^2 + 2y + (4y + 2)x + 3x^2 + x^3$.

References

- [1] R. Cori, D. Rossin, B. Salvy, Polynomial ideals for sandpiles and their Grobner bases. *Theoretical Computer Science* **276** (2002), no. 1-2, 1-15.

- [2] Chebikin Denis, Pylyavskyy Pavlo, A family of bijections between G -parking functions and spanning trees. *J. Combin. Theory Ser. A* **110** (2005), no. 1, 31–41.
- [3] Kostić Dimitrije, Yan, Catherine H. Multiparking functions, graph searching, and the Tutte polynomial. *Adv. in Appl. Math.* **40** (2008), no. 1, 73–97.
- [4] Sen-Peng Eu, Tung-Shan Fu, Symmetric parking functions and related multigraphs, *private communication*.
- [5] D. Foata, J. Riordan, Mappings of acyclic and parking functions, *Aequationes Math.* 10 (1974) 10-22.
- [6] J. Françon, Acyclic and parking functions, *J. Combin. Theory Ser. A* 18 (1975) 27-35.
- [7] J.D. Gilbey, L.H. Kalikow, Parking functions, valet functions and priority queues, *Discrete Math.* 197/198 (1999) 351-373.
- [8] Konheim, A. G. and Weiss, B. An Occupancy Discipline and Applications. *Siam Journal of Applied Mathematics* **14** (1966) 1266-1274
- [9] J.P.S. Kung, C.H. Yan, Gončarov polynomials and parking functions, *J. Combin. Theory Ser. A* 102 (2003) 16-37.
- [10] Postnikov, A. and Shapiro, B. Trees, Parking Functions, Syzygies, and Deformations of Monomial Ideals. *Transactions of the American Mathematical Society* **356** (2004).
- [11] J. Pitman, R. Stanley, A polytope related to empirical distributions, plane trees, parking functions, and the associahedron, *Discrete Comput. Geom.* 27 (4) (2002) 603-634.
- [12] J.Riordan, Ballots and trees, *J. Combin. Theory* 6 (1969) 408-411.
- [13] M.P. Schützenberger, On an enumeration problem, *J. Combin. Theory* 4 (1968) 219-221.
- [14] R.P. Stanley, Hyperplane arrangements, interval orders and trees, *Proc. Natl. Acad. Sci.* 93 (1996) 2620-2625.

- [15] R.P. Stanley, Parking functions and non-crossing partitions, in: The Wilf Festschrift, *Electron. J. Combin.* 4 (1997) R20.
- [16] C.H. Yan, Generalized tree inversions and k-parking functions, *J. Combin. Theory Ser. A* 79 (1997) 268-280.
- [17] C.H. Yan, On the enumeration of generalized parking functions, *Congr. Numer.* 147 (2000) 201-209.
- [18] C.H. Yan, Generalized parking functions, tree inversions and multicolored graphs, *Adv. in Appl. Math.* 27 (2001) 641-670.